

Statistiek I


Choice of Statistical Tests

John Nerbonne

CLCG, Rijksuniversiteit Groningen

<http://www.let.rug.nl/nerbonne/teach/Statistiek-I/>

Choice of Statistical Tests



Choice of Statistical Tests

Nominal Data

- χ^2 test of independence
 - check whether one variable influences another
 - organize one set of variable values into columns, the second into rows
 - (reformulated) question: is the distribution roughly the same in the different rows?
 - H_0 : no influence

Owner	Pet Ownership		
	Type of Pet		
	dogs	cats	other
boys	39	55	6
girls	26	65	9

- no cell with expectation of zero's
- small 2×2 tables require *Yates* correction

Nominal Data

Sign Test

- can be used, e.g.,
 - a. to test agreement in judgement
 - b. to indicate improvement
- interpretation
 - a. + indicates agreement, – disagreement
 - b. + indicates improvement, – none
- question: is the breakdown of +’s and –’s a chance breakdown?
- relatively insensitive
- almost always applicable (even when original data is not normally distributed, and even asymmetrical)

Averages

To check for a difference in averages where the standard deviation is known, the most sensitive test is the z-test.

- how many standard errors separate the two sample averages?

$$z = \frac{m_1 - m_2}{\sigma / \sqrt{n_1 + n_2}}$$

- averages are always normally distributed (no need to check for normality)
- H_0 : no difference in averages
- interpret using standard-normal tables

Averages, σ unknown

To check for a difference in averages where the standard deviation is unknown, first ask whether are testing two different groups (unrelated samples), or two scores from one group (paired data).

t-test for unrelated samples

- t statistic like z , uses s instead of σ
- H_0 : no difference in averages
- sample size is important
 - if $n \leq 15$, dist. must be normal
 - if $n \geq 40$, z is almost identical
- if t cannot be used, try **Mann-Whitney U test**
- with 3 or more groups, use **ANOVA**

Mann-Whitney U-Test

alternative to t -test (unpaired data)

- H_0 : samples from same population
- often applied to Likkert scale data
- **generalization** to several groups: **Kruskal-Wallis**

Averages, σ unknown

To check for a difference in averages where the standard deviation is unknown and two scores from one group (paired data).

t -test for paired data

- uses same t statistic as t -test for unrelated samples, uses s instead of σ
- H_0 : no difference in averages
- sample size is important
 - if $n \leq 15$, dist. must be normal
 - if $n \geq 40$, z is almost identical
- if t cannot be used, try **Wilcoxon test**

Wilcoxon

Wilcoxon's Signed Rank Test

- also applicable to ordinal data
- fallback for paired t -test
- distribution should be roughly symmetric, not skewed
- if data asymmetric, try **sign test**

Next Week—Exam

Break a leg!